

1
2
3 High quality genome **assembly** of the
4 brown hare (*Lepus europaeus*) with
5 chromosome-level scaffolding
6

7 Craig Michell^{1,2}, Joanna Collins³, Pia K. Laine⁴, Zsófia Fekete^{1,5}, Riikka
8 Tapanainen¹, Jonathan M. D. Wood³, Steffi Goffart¹ and Jaakko L. O.
9 Pohjoismäki*¹

10
11 ¹Department of Environmental and Biological Sciences, University of Eastern Finland – Joensuu, Finland

12 ²Red Sea Research Center, Biological and Environmental Science and Engineering Division, King Abdullah
13 University of Science and Technology, – Thuwal, Kingdom of Saudi Arabia

14 ³Tree of Life, Wellcome Sanger Institute, Hinxton, Cambridge, UK.

15 ⁴Institute of Biotechnology, University of Helsinki, Finland.

16 ⁵Hungarian University of Agriculture and Life Sciences, Institute of Genetics and Biotechnology, Gödöllő,
17 Hungary

18
19 *Corresponding author

20 Correspondence: jaakko.pohjoismaki@uef.fi
21

22 **Abstract**
23

24 We present here a high-quality genome assembly of the brown hare (*Lepus europaeus* Pallas),
25 based on a fibroblast cell line of a male specimen from Liperi, Eastern Finland. This brown hare
26 genome represents the first Finnish contribution to the European Reference Genome Atlas pilot
27 effort to generate reference genomes for European biodiversity.

28 The genome was assembled using 25X PacBio HiFi sequencing data and scaffolded
29 utilizing a Hi-C chromosome structure capture approach. After manual curation, the assembled
30 genome length was 2,930,972,003 bp with N50 **scaffold** of 125.8 Mb. 93.16% of the assembly
31 could be assigned to 25 identified chromosomes (23 autosomes plus X and Y), matching the
32 published karyotype. The chromosomes were numbered according to size. The genome has a
33 high degree of completeness based on the BUSCO score (mammalia_odb10 database), Complete:
34 96.1% [Single copy: 93.1%, Duplicated: 3.0%], Fragmented 0.8%, and Missing 2.9%. The
35 mitochondrial genome of the cell line was sequenced and assembled separately.

36 The brown hare genome is particularly interesting as this species readily hybridizes with
37 the mountain hare (*Lepus timidus* L.) at the species contact zone in northern Eurasia, producing

38 fertile offspring and resulting in gene flow between the two species. In addition to providing a
39 useful comparison for population studies, the genome can offer insight into the chromosomal
40 evolution among Glires in general and Lagomorpha in particular. The chromosomal assembly of
41 the genome also demonstrates that the cell line has not acquired karyotypic changes during
42 culture.

43

44 **Keywords:** Lagomorpha, reference genome, chromosome structure, phylogeny, fibroblast cell
45 line

46 Introduction

47

48 The brown hare (*Lepus europaeus* Pallas), also known as the European hare, is a widespread
49 species in the western parts of Eurasia (Bock, 2020). Besides its native range, the brown hare has
50 been introduced to numerous regions, including the British Isles, the Falkland Islands, Canada,
51 South America, Australia and New Zealand (Jaksic, 2023; Petrovan, 2013). In many places the
52 brown hare is regarded as an invasive species and a threat to the local ecosystems (Stott, 2003)
53 or native species (Reid, 2011).

54 The brown hare is a steppe-adapted species living in open grasslands and avoiding
55 forested regions. Its colonization history in Europe is complex. The species has had glacial refugia
56 during the Pleistocene in the Italian peninsula, the Balkans and Asia Minor (Fickel *et al*, 2008),
57 with an interesting pre-glacial diversity hotspot in the archipelago of Greece (Minoudi *et al*, 2018).
58 After the ice age, the species has been expanding its range both naturally as well as facilitated by
59 human-caused changes in the landscape, especially through the expansion of agricultural lands
60 and pastures. As mentioned above, the brown hare has been also frequently introduced by
61 humans to new areas from antiquity to the present day (Petrovan, 2013), resulting in potential
62 mixing of ancestral populations. This is true also for Finland, where the species has arrived
63 naturally from the southeast through the Karelian isthmus in the late 19th century (Ognev, 1940;
64 Siivonen, 1972) but also through local introductions to the southwestern parts of the country in
65 the 1910s (Rikala, 1925).

66 Curiously, Linné was unaware of the existence of the brown hare, despite having described
67 the mountain hare (*Lepus timidus* L.) and the South African Cape hare (*Lepus capensis* L.). In fact,
68 the brown hare is not native to Linné's home country Sweden but was introduced there only a
69 century after his death (Thulin, 2003). Curiously, the species author Peter Simon Pallas never
70 published a formal description of the brown hare. Instead, the species authority has been
71 attributed to him because of inclusion of the name *Lepus europaeus* into his identification table
72 of hare species of the world (Pallas, 1778: 30) (Figure 1A). Consequently, there is also no
73 information of the type locality or any other information of the species in the original publication.
74 The type locality of the brown hare has been retrospectively assigned – without too convincing
75 arguments – to Burgundy (France) or Poland (Holden, 2005). Identifying Poland as the type
76 locality was done by the Russian zoologist Sergey Ognev, and is based on Pallas mentioning in the
77 1778 text the hybrid forms of *Lepus variabilis* (synonym of *L. timidus*) and *L. europaeus* from
78 Poland and Lithuania, as well as citing later works of Pallas for the species distribution (Ognev,
79 1940). In fact, Ognev (1940: 141) boldly states that “we consider southwestern Poland as a typical

80 terrain for the nominal subspecies; there are hares with the characteristics of the basic form,
81 expressed with full clarity” (translation from Russian).

82 While being widespread in the temperate regions of Western and Central Europe to the
83 Caspian steppes in historical times, **the brown hare began a northward** expansion in the 19th
84 century. In the Fennoscandian region, the species reached the St Petersburg area by 1820s
85 (Ognev, 1940), arriving in Finland at the turn of the century (Siivonen, 1972; Thulin, 2003). By
86 1930, the species had become established throughout southern Finland, with the range
87 expansion stagnating southwest of the N64°-N62° -line until the 1990s. **Quite likely benefitting**
88 **from the ongoing anthropogenic climate change, brown hares range expansion has intensified**
89 **during the last three decades, currently reaching the polar circle (Levanen *et al*, 2018a) (Figure**
90 **1B), as well as higher altitudes in the Alps (Schai-Braun *et al*, 2023). In Finland, the expansion**
91 **seem to be limited with 150 days of annual snow cover (Levanen *et al.*, 2018a). The range**
92 expansion together with the past introductions have brought the brown hare increasingly in
93 contact with the mountain hare, especially at the northern edges of the species distribution,
94 constituting a threat especially to the more temperate climate adapted populations and
95 subspecies of the mountain hare (Reid, 2011; Reid & Montgomery, 2007; Thulin, 2003; Thulin *et*
96 *al*, 2021). Besides competing, brown hares also hybridize with mountain hares, producing fertile
97 offspring and resulting in gene flow between the species. **This gene flow is biased** towards the
98 brown hare (Ferreira *et al*, 2021; Levanen *et al.*, 2018a; Levanen *et al*, 2018b; Thulin & Tegelström,
99 2002), which obtains genetic variation from mountain hares, some of which might have adaptive
100 significance (Pohjoismaki *et al*, 2021). In contrast to the brown hare’s expansion towards north,
101 many of the Central European brown hare populations are contracting in range and numbers,
102 likely driven by changes in agricultural practices and land use (Schai-Braun *et al*, 2013; Smith *et*
103 *al*, 2005).

104 Overall, the brown hare’s features as an invasive species outside Europe and expansive
105 species in the north, its hybridization tendency and complex population ancestries make the
106 species highly interesting not only to study population genetics and adaptation mechanisms, but
107 also for the understanding the genomic makeup of species boundaries (Gaertner *et al*, 2022).
108 These studies would be greatly facilitated by the availability of high quality, chromosomally
109 scaffolded reference genomes (Blaxter *et al*, 2022).

110 **The last decade has witnessed a boom in genome sequencing technologies, leading to**
111 **many sequencing initiatives, such as the Earth BioGenome Project (EBP), European Reference**
112 **Genome Atlas (ERGA) and the Darwin Tree of Life (DTOL), aiming to produce high quality reference**
113 **genomes for a variety of organisms (Blaxter *et al.*, 2022). This has been enabled by the**
114 **technological development of high-throughput sequencing of long molecules. The read length**
115 **N50’s > 19 kbp, allows a high contiguity compared to the genome assemblies released in the early**
116 **2010’s that were based on short read sequencing technologies. Additionally, Hi-C sequencing, a**
117 **method measuring the contact frequency between all pairs of loci in the genome, has allowed**
118 **producing pretext maps of chromosomes (Ghurye *et al*, 2019), enabling chromosome level**
119 **assemblies with a precision that has been previously achieved only with model organisms**
120 **(Lawniczak *et al*, 2022).**

121 **In the presented work, we apply the PacBio HiFi long-range sequencing technology with**
122 **Hi-C sequencing to produce a high-quality reference genome of the brown hare (*Lepus***
123 ***europeus*). While there is no existing genome assembly for the species, a mountain hare**

124 assembly (GCA_009760805) exists, representing the Irish subspecies *L. t. hibernicus* and is based
125 on a female specimen, thus lacking the Y-chromosome. This assembly is also a so-called
126 “pseudoreference”, assembled to chromosomes using the rabbit (*Oryctolagus cuniculus* L.)
127 reference genome (Marques *et al.*, 2020). The original rabbit reference genome had been
128 established using whole-genome shotgun sequencing of females representing the inbred
129 Thorbecke New Zealand White line and was quite fragmented (Carneiro *et al.*, 2014), updated only
130 recently with long-read sequencing data (Bai *et al.*, 2021). Notably, the diploid chromosome
131 number in domestic rabbits is $2n = 44$ (Korstanje *et al.*, 1999), whereas it is 48 in brown hare and
132 mountain hare (Gustavsson, 1972).

133 To obtain a reference genome conforming to current standards, we used a fibroblast cell
134 line (LE1) of a male specimen of brown hare (*Lepus europaeus*) from Liperi, Eastern Finland
135 (Gaertner *et al.*, 2022), as a source of high-molecular weight (HMW) DNA and fresh RNA.
136 Compared to solid tissues, fibroblasts are optimal for Hi-C, as the method was originally
137 developed for cells growing in a monolayer (Lieberman-Aiden *et al.*, 2009). Now, over 240 years
138 after the first mention of the name *Lepus europaeus* in the literature (Figure 1A), we have
139 expanded the original seven-word description to encompass 2,930,972,003 genomic letters.
140 These letters have been assembled in a highly complete (BUSCO mammalia_odb10 database,
141 complete: 96.1%), and continuous (N50 scaffold: 125.8 Mb) manner with 93.16% of the sequence
142 being placed in the identified 23 autosomes, and X and Y sex chromosomes. The presented
143 genome assembly not only contains the identity of the brown hare as a species, but also provides
144 a gateway to detailed knowledge of its biology and evolutionary history. Besides being the first
145 Finnish contribution to the European Reference Genome Atlas (Mazzoni *et al.*, 2023), the brown
146 hare reference genome represents a continuing effort to map and understand our planets
147 biodiversity.

148 **Methods**

149

150 **Sampling and confirming of the species identity**

151 A young male brown hare was hunted in October 2018 in Kuorinka, Liperi, Finland
152 (62.6207 N, 29.4478 E, Figure 1B). In Finland, brown hares are highly dependent on similar man-
153 made environments (Levanen *et al.*, 2018a; Levänen *et al.*, 2019), and also the collection location
154 is agricultural area with a mosaic of fields, shrubs and mixed forest with a strong brown hare
155 population and only occasional stray mountain hares. Geographically, the sampled population is
156 part of the same distribution continuum as with species proposed type locality, through Russian
157 Karelia and the Baltic states to Poland. This is important, as is recommendable that the species
158 reference genome would represent or be closely related to the type locality population
159 (Lawniczak *et al.*, 2022).

160 The sampling had minimal impact on the local brown hare population and no impact on
161 the habitat. As brown hares are legal game animals in Finland and the hunting followed the
162 regional hunting seasons and legislation (Metsästyslaki [Hunting law] 1993/615/5§), the sampling
163 adheres to the ARRIVE guidelines and no ethical assessment for the sampling was required. The
164 sampling also did not involve International Trade in Endangered Species of Wild Fauna and Flora
165 (CITES) or other export of specimens, as defined by the Convention on Biological Diversity (CBD).

166 The species identity was confirmed at site based on the morphological features
167 distinguishing the brown hare from the mountain hare, the only other hare species in northern
168 Europe. Further analysis of the coding part of the LE1 genome and mitochondrial DNA
169 haplotyping showed minimal ancestral admixture with **mountain hares (Gaertner *et al.*, 2022)**. **In**
170 **fact, unlike many other brown hares in the region, showing adaptive introgression of mountain**
171 **hare specific UCP1 alleles (Pohjoismaki *et al.*, 2021), the LE1 specimen is homozygous for the**
172 **common ancestral brown hare allele, UCP02.**

173

174 **Generation and vouchering of the cell line**

175 The fibroblast cell line (LE1) was isolated from the specimen as described earlier (Gaertner
176 *et al.*, 2022) and is deposited as cryopreserved living cells under voucher number ZFMK-TIS-69747
177 into the biobank of the Stiftung Leibniz-Institut zur Analyse des Biodiversitätswandels, Zoological
178 Research Museum Alexander König (ZFMK), Bonn, Germany. Additional identifiers for the sample
179 are mLepEur2 and ERGA specimen ID ERGA_FI_3610_002 (COPO portal, [https://copo-](https://copo-project.org)
180 [project.org](https://copo-project.org)).

181

182 **HMW DNA extraction and PacBio HiFi sequencing**

183 A salting out-method was used to extract high molecular weight DNA from cells grown to
184 confluency on a 10 cm cell culture dish. Briefly, the cells were detached from the dish using
185 trypsin, followed by centrifugation and two washes with PBS. The salting out-method then
186 followed the 10X Genomics “Salting Out Method for DNA extraction from cells” protocol (**10x-**
187 **Genomics, 2017)**. The high molecular weight DNA was quantified using a Qubit 3.0 (**Thermo Fisher**
188 **Scientific™, Malaysia)**, followed by qualification with an 0.8 % Agarose gel. The DNA was **sent for**
189 **the library preparation and sequencing at the** DNA Sequencing and Genomics Laboratory,
190 Institute of Biotechnology, University of Helsinki on the PacBio Sequel II. **In brief, the DNA was**
191 **quantified using Qubit and DNA fragment length was measured using Fragment Analyzer (HS large**
192 **fragment kit). DNA was then sheared with Megaruptor 3 (Diagenode) (45/200/30) to produce**
193 **DNA of length 24 kb. Buffer was then replaced with PacBio’s Elution buffer using AMPure beads.**
194 **Repair and A-tailing, adapter ligation, nuclease treatment and cleanup with SMRTbell cleanup**
195 **beads was then performed with the SMRTbell prep kit 3.0. Larger fragments (greater than 10 000**
196 **bp) were then purified with the BluePippin (Sage Science) (0.75% DF Marker S1 high pass 6-10kb**
197 **v3) and DNA was further purified with PacBio AMPure beads and treated with DNA damage repair**
198 **mix (PacBio) (37oC for 60 mins) and then purified with AMPure clean-up beads and eluted to 11**
199 **ul PacBio’s Elution buffer. Libraries were sequenced on the PacBio Sequel II at a concentration of**
200 **90 pM using PacBio’s instructions (provided by the SMRTlink software). Sequence data from two**
201 **flow cells was produced. Genomescope2 was used to visualize the HiFi sequencing results**
202 **(Ranallo-Benavidez *et al.*, 2020).**

203

204 **Mitochondrial DNA (mtDNA) sequencing**

205 The mitochondrial genomes of our hare cell lines (Gaertner *et al.*, 2022) have been
206 sequenced for the purposes of another study using approximately 2 kb overlapping PCR-amplified
207 fragments of mtDNA. The primers used to amplify the mountain hare mtDNA were as follows:

208 Le93F: TTGTTTTGTAGCAAGTTTACACATGC

209 Le184R: GCTTAATACCTGCTCCTCTTGATCTA

210 Le1580F: TTAAACCCATAGTTGGCCTAAAAGC
211 Le1635R: TTGAGCTTTAACGCTTTCTTAATTGA
212 Le3045F: AGGCGTATTATTTATCCTAGCAACCT
213 Le3175R: CCTCATAAGAAATGGTCTGTGCGA
214 Le3921F: CCCCTAATCTTTTCCATCATCCTAT
215 Le4482R: TCATCCTATATGGGCAATTGAGGAAT
216 Le4689F: AGGCTTTATTCCAAAGTGAATTATTATTCA
217 Le5417R: AGGCTCCAAATAAAAGGTAGAGAGTT
218 Le6696F: ATACCGTCTCATCAATAGGCTCCTTC
219 Le6756R: ATAAAGATTATTACTATTACAGCGGTTAGA
220 Le8603F: AGCCTATATCTACATGATAATACTTAATGA
221 Le8698R: CGGATAAGGCCCCCGTAAGTGG
222 Le10552F: TTGAAGCAACACTAATCCCTACACTA
223 Le10613R: TCGTTCTGTTTGATTACCTCATCGT
224 Le11301F: ACCATTAACCTTCTAGGAGAGCTTCT
225 Le11807R: AGGATAATGATTGAGACGGCTATTGA
226 Le12407F: GTCTAATCCTAGCTGCTACAGGTAAG
227 Le12791R: GAGCATAAAAAGAGTATAGCTTTGAA
228 Le14204F: ATTGTTAACCACTCTCTAATCGACCT
229 Le14514R: CCAATGTTTCAGGTTTCTAGGTAAGT
230 Lt16056F: TGGGGTATGCTTGGACTCAAC
231 Le16119R: TCGTCTACAATAAGTGCACCGG
232

233 In total, 12 separate reactions were prepared to cover the mitochondria genome:


- 234 1. Lt16056F + Le184R: 1871 bp
- 235 2. Le93F + Le1635R: 1543 bp
- 236 3. Le1580F + Le3175R: 1596 bp
- 237 4. Le3045F + Le4482R: 1438 bp
- 238 5. Le3921F + Le5417R: 1497 bp
- 239 6. Le4689F + Le6756R: 2068 bp
- 240 7. Le6696F + Le8698R: 2003 bp
- 241 8. Le8603F + Le10613R: 2011 bp
- 242 9. Le10552F + Le11807R: 1256 bp
- 243 10. Le11301F + Le12791R: 1491 bp
- 244 11. Le12407F + Le14514R: 2108 bp
- 245 12. Le14204F + Le16119R: 1916 bp

246 (Expected fragment size based on the published *Lepus europaeus* mtDNA sequence from Sweden
247 [NC_004028.1]).
248

249 The fragments were amplified from total DNA preparations using a PCR program with a 1 min
250 94 °C denaturing step, followed by 35 cycles of 94 °C for 15 s, 56 °C for 15 s and 72 °C for 2 min
251 and a final 3 min elongation step at 72 °C. The obtained products were gel purified using the
252 GeneJET gel extraction kit (Thermo Fisher Scientific™) and sent for sequencing using Illumina
253 MiniSeq at the Genome Center of Eastern Finland.

254
255
256
257
258

The sequence of the non-coding region-containing PCR fragment (Lt16056F + Lt184R) was further validated by Sanger sequencing, applying also the following additional sequencing primer: Le101F: TATAAATTCCTGCCAAACCCCAAAAA



259
260
261
262
263
264
265
266
267
268
269

Figure 1. The brown hare. (A) An excerpt from the identification table by Pallas (1778: 30), which has been recognized as the species description for *Lepus europaeus*. “Tips of the ears longer than the head, tail black above.” (translation from Latin). (B) The geographic location of the LE1 sample used in this study. The grey-shaded area shows the approximate current distribution of the brown hare in Finland, based on the data in the Finnish Biodiversity Information Facility (FinBIF, <https://laji.fi/en>). Individual records from the north are from settlements or towns. The arctic circle (black line) runs along the 66.6 ° parallel. Background photo shows a typical Finnish male brown hare. Photo taken in Joensuu, 20 km East from the LE1 sampling site by Dr. Mervi Kunnasranta.

270 Hi-C library preparation

271 Hi-C sequencing libraries were prepared following the protocol of (Belaghzal *et al*, 2017)
272 with the following changes: 1.) To prepare a diverse sequencing library, we performed the Hi-C
273 protocol in triplicates, 2.) Size fractionation was performed using the NEBNext Ultra II FS DNA
274 module (New England Biolabs), 3.) Library preparation was performed using the NEBNext Ultra II
275 library preparation kit for Illumina (New England Biolabs) and 4.) Library enrichment was

276 performed using triplicate PCR reactions with six cycles of PCR. The PCR reactions were then
277 purified using Ampure XP beads (Beckman Coulter Life Sciences) at a ratio of 0.9X. The final clean
278 libraries were quantified using Qbit, followed by agarose gel electrophoresis to confirm the
279 fragment size. The sequencing was performed on a single lane of the Illumina NovaSeq 6000 using
280 the SP flowcell with paired-end chemistry 2 x 150bp.

281

282 **Genome assembly**

283 HiFiasm version 0.18.7 (Cheng *et al*, 2021) was used to assemble the 25X (rq >0.99) PacBio
284 HiFi reads using the recommended arguments *-l3 -h1 and -h2 -primary* to integrate the Hi-C read
285 data and produce a primary assembly. We then continued with the scaffolding of the primary
286 assembly and annotation. To process the HiC data, we first mapped the Hi-C data to the primary
287 genome assembly using BWA-mem version 0.7.17 (Li & Durbin, 2009) with the arguments *-SP*.
288 The mapped reads were then parsed and filtered using pairtools version 1.0.2 (Open2C *et al*,
289 2023). To parse the aligned Hi-C reads, we used the options *-no-flip -dd-columns mapq -walks-*
290 *policy mask*. The parsed pair data was then sorted and deduplicated using default arguments *with*
291 *pairtools*. Finally, we selected unique-unique and unique-rescued pairs and split these into the
292 pairs file and bam file for input in YaHS version 1.1 (Zhou *et al*, 2023). YAHS was run using the
293 default parameters with the primary contig assembly and the filtered Hi-C bam file. **HiFiasm**
294 **operates as a haplotype-aware assembler, counting and assembling unique k-mers. Utilizing k-**
295 **mer coverage information, the assembler identifies regions in the genome with expected**
296 **heterozygous coverage and subsequently amalgamates them into a unified sequence. To refine**
297 **the assembly further, Hi-C data is incorporated to elucidate the phasing of variants and ascertain**
298 **connected genomic regions. Consequently, the assembler's output provides a primary assembly**
299 **along with an alternative assembly, effectively separating the haplotypes by their distinct alleles**
300 **within the genomic content (please see [https://lh3.github.io/2021/04/17/concepts-in-phased-](https://lh3.github.io/2021/04/17/concepts-in-phased-assemblies)**
301 **[assemblies](https://lh3.github.io/2021/04/17/concepts-in-phased-assemblies)).** Contiguity and general genome statistics were calculated using QUASt version 5.2.0
302 (Mikheenko *et al*, 2018). We assessed the completeness of the genome by calculating the number
303 of complete single copy orthologs with BUSCO version 5.1.2 (Manni *et al*, 2021), using the
304 *mammalia_odb10* database as well as the more lineage specific *glires_odb10* database.

305

306 **Genome annotation**

307 Repeat annotation of the genome was performed with EDTA version 2.1.0 (Ou *et al*, 2019),
308 a *de novo* repeat identification pipeline. Using the repeat library produced by EDTA, we masked
309 the scaffolded genome using RepeatMasker version 4.1.1 (Smit *et al*, 2013-2015). RNA-seq data
310 from the same cell line (SRA accession number: SRR18740842) as well as RNA-seq data from other
311 *L. timidus* libraries (SRA accession number: SRR10020054, SRR10020055, SRR10020060,
312 SRR10491719, SRR18740839, SRR18740840, and SRR18740841) was collected from the sequence
313 read archive (SRA). **The previously produced RNA-seq data (Gaertner *et al.*, 2022)** was trimmed
314 using fastp version 0.23.2 (Chen *et al*, 2018) and mapped against the masked genome using
315 HISAT2 version 2.1.0 (Kim *et al*, 2019) with the default parameters. Furthermore, we included the
316 protein sequences of the assembled transcripts from this cell line as further evidence for the
317 genome annotation. These lines of gene evidence were included in the annotation using BRAKER3
318 version 3.02 (Bruna *et al*, 2021). Telomeric sequences [AACCCT]*n* were identified using a Telomere

319 Identification toolkit (tidk) version 0.2.31. The telomeric sequence copy number was then
320 calculated in windows of 200kb for visualization using Circos (Krzywinski *et al*, 2009).

321

322 **Mitochondrial DNA assembly and annotation**

323 Mitochondrial DNA was assembled from the PCR-amplified and Illumina sequenced
324 mtDNA using the MitoZ pipeline (Meng *et al*, 2019). After comparing the results of the pipeline's
325 outputs with different kmer options, we selected the best assembly as final. Run options used in
326 the final assembly were --clade Chordata --fastq_read_length 150, --requiring_taxa Chordata --
327 genetic_code 2 --kmers_megahit 21 29 39 59 79 99 119 141. The tools invoked by the pipeline
328 included fastp (Chen *et al.*, 2018) for cleaning the raw data, MEGAHIT (Li *et al*, 2015) for assembly,
329 after which sequences were filtered to ensure the correct taxa by HMMER (Wheeler & Eddy, 2013)
330 and further filtered for completeness of protein-coding genes. Annotation steps were done using
331 TBLASTN (Gertz *et al*, 2006), GeneWise (Birney *et al*, 2004) and MiTFi (Juhling *et al*, 2012). Final
332 manual curation, the annotation of the non-coding region (NCR) as well as the illustration of the
333 mitochondrial genome was done using Geneious® 10.2.6 (Biomatters. Available from
334 <https://www.geneious.com>). The functional loci on the NCR were identified based on the
335 similarity with the human (NC_012920) and mouse (FJ374652) NCR sequences.

336

337 **Manual curation**

338 The assembled and annotated genome was manually curated to further improve its
339 quality based on Hi-C contact maps as described in (Howe *et al*, 2021), allowing to break or join
340 erroneous scaffold assemblies and removing duplicated contigs.

341

342 **Comparison with previous assembly**

343 We performed a comparison of our scaffolded assembly to the current *L. timidus* genome
344 assembly (NCBI Accession number: GCA_009760805). Mapping to the genome was performed
345 using minimap2 version 2.21 (Li, 2018) with the arguments *-asm5*. A dot plot of the alignment
346 was created using the R script pafCoordsDotPlotly.R (<https://github.com/tpoorten/dotPlotly>).

347

348 **Results**

349

350 The genome assemblies can be accessed via BioSample accession SAMN33984520 as well as
351 BioProject accession PRJNA1009711 for the primary assembly and PRJNA1009710 for the
352 alternative assembly.


353

354 **Genome assembly**

355 The expected haploid genome size of the *L. europaeus* is similar to *L. timidus*, which has
356 been estimated to be 3.25 pg (3,178,500,000 bp) (Vinogradov, 1998), containing 23 autosomal
357 chromosomes and two sex chromosomes (Gustavsson, 1972). PacBio HiFi sequencing with two
358 flow cells resulted in 25 X coverage of the expected genome size. The sequence N50 of the HiFi
359 data was 19.97 kb. When k-mer size of 21 is used, the resulting genome size from the PacBio HiFi
360 data is 2.4 Gbp (Figure 2A). The Illumina sequencing of the Hi-C data produced 494,285,585 paired
361 reads, representing about 51 X coverage of the genome, with a duplication rate of 17 %. Assembly

362 with HiFiasm yielded a contig assembly of 2.96 Gbp made up of 671 contigs with a contig N50 of
363 58 Mbp. The longest contig was 149 Mbp. Using the uniquely mapped Hi-C data, we were able to
364 scaffold the contigs and fix misassembled contigs (Figure 2B). **Although the genome size (2.96**
365 **Gbp) and the largest scaffold (149 Mbp) were similar to the contig assembly, the scaffold N50 was**
366 **increased to 124 Mbp (N=10) after the Hi-C scaffolding. The obtained scaffold N90 was 21 Mbp**
367 **(N=25).** The BUSCO scores of the *L. europaeus* assembly suggest it to be near-complete, with the
368 following results (%- of the total genome): Complete: 96.1% [Single copy: 93.1 %, Duplicated: 3.0
369 %], Fragmented 0.8 %, and Missing 2.9 % based on the mammalia_odb10 database.

370 To further improve the assembly, manual curation was performed, resulting in 76 scaffold
371 breaks, 81 joins and removal of 62 erroneously duplicated contigs (Table 1). These interventions
372 led to an increase in scaffold N50 of 1.69 % from 123.7 Mb to 125.8 Mb, and a reduction in scaffold
373 count of 9.21 % from 716 to 650. Of the finalized assembly, 93.16 % could be assigned to 25
374 identified chromosomes (23 autosomes plus X and Y), matching the known karyotype.
375 Chromosomes were numbered according to size (Table 2). The curated genome length was
376 2,930,972,003 bp.



377
 378 **Figure 2.** A) Genomescope2 profile of the PacBio HiFi data. B) YaHS pretext map of Hi-C
 379 scaffolding after manual curation. C) Dot plot comparison with the mountain hare
 380 pseudoreference (GCA_009760805) assembly with rabbit chromosomal assignments.
 381 Each grey horizontal line represents a contig end. Names of the non-chromosomal
 382 contigs have been omitted for clarity. Rabbit chromosomes corresponding two hare
 383 chromosomes as well as the X chromosome with an inversion are highlighted.
 384
 385

Table 1. The brown hare reference genome chromosome assignment and assembly statistics

Chromosome assignment	
Chr length	2,730,543,680 bp
Autosomes	23
X	1
Y	1
Unlocalised	19

Statistics for primary assembly after manual curation	
# scaffolds	650
Total scaffold length	2,930,972,003
Average scaffold length	4,509,187.70
Scaffold N50	125,776,599
Scaffold auN	117,151,857.99
Scaffold L50	10
Largest scaffold	184,151,585
Smallest scaffold	1000
# contigs	1068
Total contig length	2,930,889,045
Average contig length	2,744,278.13
Contig N50	43,409,466
Contig auN	47,572,960.95
Contig L50	21
Largest contig	141,329,000
Smallest contig	1000
# gaps in scaffolds	418
Total gap length in scaffolds	82,958
Average gap length in scaffolds	198.46
Gap N50 in scaffolds	200
Gap auN in scaffolds	199.70
Gap L50 in scaffolds	208
Largest gap in scaffolds	200
Smallest gap in scaffolds	31
Base composition	
A	821,587,687
C	645,897,412
G	644,604,632
T	818,799,314

GC content	44.03 %
# soft-masked bases	0
# segments	1,068
Total segment length	2,930,889,045
Average segment length	2,744,278.13
# gaps	418
# paths	650

387
388
389

Table 2. Chromosome assemblies of the brown hare reference genome with their corresponding GenBank access numbers.

Chromosome	GenBank	Size (bp)	GC content (%)
1	CM065551.1	184,151,585	42.0
2	CM065552.1	176,467,698	42.0
3	CM065553.1	169,342,607	42.0
4	CM065554.1	166,606,208	43.0
5	CM065555.1	159,335,674	46.0
6	CM065556.1	132,443,831	42.0
7	CM065557.1	132,293,278	44.0
8	CM065558.1	130,587,543	41.0
9	CM065559.1	125,776,599	44.5
10	CM065560.1	125,704,418	46.0
11	CM065561.1	117,559,561	44.0
12	CM065562.1	105,667,338	44.0
13	CM065563.1	100,925,866	44.0
14	CM065564.1	99,301,477	44.0
15	CM065565.1	95,895,056	42.0
16	CM065566.1	91,637,918	42.5
17	CM065567.1	84,898,765	45.5
18	CM065568.1	77,563,730	48.5
19	CM065569.1	72,700,603	47.5
20	CM065570.1	69,331,453	45.5
21	CM065571.1	60,609,710	47.5
22	CM065572.1	44,025,767	47.0
23	CM065573.1	30,716,498	53.0
X	CM065574.1	135,183,192	40.5
Y	CM065575.1	33,560,193	41.0
MT	OR915849.1	17,540	38.6

390

391 **Genome annotation**

392 EDTA produced a curated custom repeat library of 7,045 repetitive elements. Interestingly,
393 the proportion of the genome masked as repetitive elements was higher than expected at 46.8 %
394 of the genome. In the mountain hare **pseudoreference genome assembly (GCA_009760805)**, only
395 28 % of the genome was masked, and similarly the *k*-mer based estimate of the repetitive
396 elements was 26.7 %. To annotate the genome, we mapped ~1.6 billion RNA-seq reads from
397 previously published RNA-seq libraries, including this cell line. The average mapping rate of all
398 libraries was 89.95 %, which we consider a good indicator of the quality of this genome. Using
399 BRAKER3, we were able to annotate 19,906 gene models in the genome. We then ran BUSCO on
400 the predicted genes and the results show a good level of completeness: Complete: 81.1 % [Single
401 copy: 63.3 %, Duplicate: 17.8 %], Fragmented: 1.1 %, Missing: 17.8 %, Total: 9226 when compared
402 to the mammalia_odb10 database. Telomeric sequences on both ends of the chromosome were
403 detected on 13 chromosomes (Figure 3A). Furthermore, telomeric sequences were found in high
404 copy number throughout the chromosome length, a feature previously noted using FISH in
405 mountain hares (Forsyth *et al*, 2005).

406

407 **Comparison to the mountain hare assembly**

408 Minimap2 was able to align 97.3 % of the contig sequences from the mountain hare
409 **pseudoreference genome assembly (GCA_009760805)** to this genome **assembly** (Figure 2C). **As**
410 **pointed out earlier, the mountain hare genome assembly has been scaffolded against the rabbit**
411 **reference genome (Giska *et al*, 2022), which has a different chromosome count (Beklemisheva *et***
412 ***al*, 2011) and explains the few notable inversions and structural differences between the two**
413 **genomes (Figure 2C). For example, the chr7 and chr 12 assemblies of the brown hare have a good**
414 **degree of synteny with the mountain hare pseudoreference (= rabbit) chr1, with the chr12 being**
415 **in inverted orientation. Similarly, the brown hare chr13 and chr16 assemblies have a good synteny**
416 **and orientation with the rabbit chr2.**

417

418 **Mitochondrial genome**


419 While it is possible to recover the entire mitochondrial genome from the HiFi sequencing
420 data (Uliano-Silva *et al*, 2023), we had already previously sequenced the mitochondrial genomes
421 of our hare cell lines for the purposes of another study. The assembly of the Illumina data
422 provided a 16,836 bp mtDNA sequence, which was complete and circular, with a 142 bp non-
423 repetitive overlap between the 3' and 5' ends of the original assembly removed from the final
424 version. All the expected vertebrate mitochondrial genes, 37 in total, were found (Figure 3B).

425 Interestingly, the primary assembly of the Illumina data was notably smaller than
426 expected based on the original PCR amplicons, with a notable difference between the expected
427 and obtained non-coding region (NCR) length. Sanger sequencing of the PCR-fragment containing
428 the NCR (Lt16056F + Le184R) revealed that it contained a sequence run consisting of six head-to-
429 tail repeats of a 140 bp element between OriH and tRNA-Phe, explaining the incorrect assembly
430 of the NCR sequence from the short-read Illumina data. With the addition of the repeat elements,
431 the final mitochondrial genome size of our specimen is 17,540 bp.

432 Based on the sequence similarity with human and mouse mitochondrial genomes, we
433 were able to tentatively identify the termination associated sequence (TAS, 2 nt difference to
434 human), required for the generation of the 7S DNA and thus the displacement loop (D-loop)

435 structure on the NCR (Crews *et al*, 1979). Interestingly, assuming that the 7S DNA is roughly the
 436 same size as in other mammals (600-650 nt), its 5'-end would map to the beginning of another
 437 highly repetitive 240 bp region, likely to have structural significance. This is a notably longer
 438 sequence run than the tRNA-like sequence reported at the replication origin of *L. timidus* mtDNA
 439 (Melo-Ferreira *et al*, 2014). We have preliminary assigned this structure to correspond to the
 440 heavy-strand origin of replication (OriH), supported by the fact that a similar, tRNA sequence-
 441 derived hairpin structure is required for the priming of the origin of the light-strand replication
 442 (OriL) (Fuste *et al*, 2010). Experimental validation of the proposed OriH is required.

443
 444



445 **Figure 3.** A reference genome for the brown hare. A) A size map of 23 autosomes as well as X and
 446 Y sex chromosomes of *Lepus europaeus*. Telomeric sequences and their relative lengths are
 447 indicated on the inner circle. B) A schematic illustration of the mountain hare mitochondrial
 448 genome. Genes encoded by the H-strand are placed inside and the L-strand genes outside of the
 449 circle. NCR = non-coding region.

451 Discussion

452
 453 Given the challenges of assembling a large mammalian genome, we have produced a high-
 454 quality genome **assembly** for the brown hare. The genome **assembly** has a high degree of
 455 contiguity (Contig N50 43 Mbp) and completeness (BUSCO complete 96.1 %). Interestingly, the
 456 genome contains a high number of repetitive elements, a fact that warrants further investigation
 457 to elucidate their identity. **While vast majority of the sequences seem to come from**
 458 **retroelements or endoretroviruses, we cannot fully exclude that some multicopy host genes could**
 459 **be masked. A detailed analysis of the repeat element libraries from this and another vertebrate**
 460 **genome assembly releases might be warranted.**

461 The genome **assembly** is in chromosome scale, and all 23 autosomes as well as X and Y
 462 chromosomes could be scaffolded. Compared with **the closely related mountain hare**

463 pseudoreference assembly, the brown hare genome assembly is much more contiguous (Table 1)
464 and inclusion of telomeric repeats on both ends of several chromosomes (Figure 3A) further
465 demonstrates the telomere-to-telomere continuity of these assemblies. The presented assembly
466 also resolves the expected chromosome structure for the species (Figure 2B, C), which is
467 informative regarding the karyotype difference between hares ($n = 24$) and rabbits ($n = 22$).
468 Comparison of the brown hare assembly with the mountain hare pseudoreference chromosomes
469 shows that the rabbit chromosome 1 corresponds to the brown hare chromosomes 12 and 17,
470 with the chromosomal arm corresponding to the former is in an inversed orientation. Similarly,
471 the rabbit chromosome 2 corresponds to the hare chromosomes 13 and 16, both in the same
472 orientation. It should also be noted that as the hare chromosomes were numbered according to
473 their size, the numbering in general does not match those of the rabbit chromosomes (Figure 2C)

474 The obtained 17,540 bp long mitochondrial genome assembly also provides some
475 interesting aspects of mtDNA sequence variation within mammalian species. The size difference
476 to human (16,569 bp) and mouse (16,298 bp) mitochondrial genomes is caused by a rather long
477 non-coding region (2,102 bp vs. 1,123 bp and 879 bp in humans and mice, respectively), longer
478 rRNA and tRNA genes, as well as additional non-coding nucleotides between genes. The
479 mitochondrial genome of our specimen is slightly smaller than the 17,734 bp previous NCBI
480 Reference Sequence for brown hare (NC_004028.1) from Sweden. The size difference is mostly
481 due to a copy number difference in tandem repeat elements within the non-coding region, with
482 seven in the NC_004028.1 and six in our specimen. Indels elsewhere in the NCR explain the rest
483 of the difference.

484 The main technical challenge for highly continuous, good quality reference genome
485 assemblies is the availability of relatively large quantities of intact HMW DNA for HiFi sequencing.
486 This typically requires snap freezing of tissue samples and assuring an intact cold chain of below
487 $-80\text{ }^{\circ}\text{C}$ for their sending, handling, and long-term storage (Januszczak & Holroyd, 2023). While
488 development in sequencing technologies can reduce the required DNA amounts, obtaining high
489 sequencing coverage over a large vertebrate genome is easier when material is plentiful. Similarly,
490 RNA is very sensitive to degradation in post-mortem tissue samples, also requiring immediate
491 preparation or snap-freezing to be suitable for RNA-seq. Although transcript data is not obligatory,
492 it is highly useful for the purpose of genome annotation. The requirement of large samples of
493 fresh tissue for -omics purposes complicates the sampling of most vertebrates species. For
494 example, in Finland all land vertebrates are protected by law (animal protection law 247/1996,
495 hunting law 615/1993), with very few exceptions, such as game animals and pest species.
496 Capturing and lethal sampling is especially problematic for endangered species. Not only will the
497 sampling require extensive permits, but it is also difficult to justify and conduct ethically. Here,
498 the generation of primary fibroblast cell lines can offer a solution. Although the source specimen
499 for the cell line utilized in this study was hunted, fibroblasts can be isolated from relatively small
500 skin biopsies, such as ear clippings (Seluanov *et al*, 2010), with minimally invasive sampling and
501 harm to the individual. These cells can be expanded in tissue culture and, when immortalized,
502 provide a highly scalable source of fresh DNA and RNA. Furthermore, living cells can be stored as
503 frozen stocks in biobanks for decades (Fazekas *et al*, 2017). Tissue sampling can be done directly
504 into the growth medium in ambient temperature, allowing several days for the transport to the
505 laboratory without a need for dry ice or N_2 -cryocontainers, which greatly simplifies collection and
506 logistics. The same applies to the isolated cells. As an example, we sent living cell suspensions

507 from Joensuu, Finland to the ZFMK biobank in Bonn, Germany, over regular airmail in ambient
508 temperature. The receiving laboratory can then amplify the cells by culturing and cryopreserve
509 them in sufficient replicate stocks. It is worth to point out that the obtained genome **assembly**
510 scaffolded to 23 autosomes, X and Y chromosomes, as expected for the species, shows no
511 evidence of karyotypic drift caused by cell culture, dispelling possible concerns for the use of cell
512 lines for such work (Gaertner *et al.*, 2022).

513 The brown hare is an iconic European mammal, locally familiar to a wider public of nature-
514 goers, farmers and hunters, while being at the same time an invasive species on many other
515 continents. It has populations with complex ancestral makeup due to isolated Pleistocene refugia,
516 frequent introductions and cross-species hybridization, making it an exceptionally interesting
517 mammal for genetic studies. **Furthermore, depending on the local environmental context, brown**
518 **hares can be both an endangered as well as a rapidly expanding species, prompting opportunities**
519 **for both conservation as well as invasion genetics studies.** A high-quality reference genome
520 **assembly** will allow us also to peer deeper into genome structure and chromosome evolution,
521 which will be particularly interesting from the viewpoint of genetic introgression (Juric *et al.*, 2016)
522 and maintenance of species boundaries. Any issues of genetic compatibility between species
523 should leave their fingerprint in the genomes (Skov *et al.*, 2023), providing interesting insight into
524 the speciation mechanisms (Wolf *et al.*, 2010). Having a high-quality genome for both species has
525 important implications for the fields of evolutionary and conservation biology. Due to the high
526 contiguity of the genome **assembly**, it will be possible to perform accurate analyses of linkage
527 disequilibrium as well as the identification of runs of homozygosity (ROH). Accurate ROH analysis
528 is useful for identifying and understanding the impact of inbreeding burden on the genome of
529 increasingly threatened and isolated species, as quantitative traits can be associated with ROH
530 burden (Ceballos *et al.*, 2018). Furthermore, this information might provide important insights
531 into the hybridization of the brown hare with the mountain hare, as well as help to pinpoint
532 genomic regions that might be helping the species to adapt to local environmental conditions
533 during the ongoing range expansion (Pohjoismaki *et al.*, 2021). Other *Lepus* reference genome
534 **assemblies** available soon will help to complete this picture. Together, this genome **assembly** and
535 the mountain hare genome **assembly** will provide a solid base for future studies.

536

537 **Acknowledgements**

538 We acknowledge the DNA Sequencing and Genomics Laboratory, Institute of Biotechnology,
539 University of Helsinki for the PacBio and Hi-C sequencing, with special thanks to Dr Petri Auvinen
540 and Dr Martyn James for helpful discussions and personal interest towards the project. Dr Seppo
541 Helisalme and Dr Joose Raivo from the Gene diagnostics lab at Genome Center of Eastern Finland
542 are thanked for providing the mtDNA sequencing service. We are grateful to the European
543 Reference Genome Atlas (ERGA) for the opportunity to include our mountain hare genome into
544 their pilot project. Dr Ann McCartney is thanked for her kind help throughout the process and Mr
545 Jukka Pusa for his contribution in obtaining the brown hare specimen. We thank Ms Anita
546 Kervinen for her valuable laboratory assistance and Dr. Mervi Kunnasranta for kindly allowing to
547 use her hare photo in the publication.

548 **Data, scripts, code, and supplementary information availability**

549 The genome assembly is available from the NCBI database under BioProject ID PRJNA950335. The
550 cell line is available through the corresponding author as well as from the ZFMK biobank.

551

552 **Conflict of interest disclosure**

553 The authors declare that they comply with the PCI rule of having no financial conflicts of interest
554 in relation to the content of the article.

555

556 **Funding**

557 This study belongs to the xHARES consortium funded by the R'Life initiative of the Academy of
558 Finland, grant no. 329264.

559

560 **References**

561

562 10x-Genomics (2017) Salting out method for DNA extraction from cells. *Sample Preparation*
563 *Demonstrated Protocol* CG00016. Rev A.: 1-5

564 Bai Y, Lin W, Xu J, Song J, Yang D, Chen YE, Li L, Li Y, Wang Z, Zhang J (2021) Improving the genome
565 assembly of rabbits with long-read sequencing. *Genomics* 113: 3216-3223

566 Beklemisheva VR, Romanenko SA, Biltueva LS, Trifonov VA, Vorobieva NV, Serdukova NA,
567 Rubtsova NV, Brandler OV, O'Brien PC, Yang F *et al* (2011) Reconstruction of karyotype evolution
568 in core Glires. I. The genome homology revealed by comparative chromosome painting.
569 *Chromosome Res* 19: 549-565

570 Belaghzal H, Dekker J, Gibcus JH (2017) Hi-C 2.0: An optimized Hi-C procedure for high-resolution
571 genome-wide mapping of chromosome conformation. *Methods* 123: 56-65

572 Birney E, Clamp M, Durbin R (2004) GeneWise and Genomewise. *Genome Res* 14: 988-995

573 Blaxter M, Archibald JM, Childers AK, Coddington JA, Crandall KA, Di Palma F, Durbin R, Edwards
574 SV, Graves JAM, Hackett KJ *et al* (2022) Why sequence all eukaryotes? *P Natl Acad Sci USA* 119

575 Bock A (2020) *Lepus europaeus* (Lagomorpha: Leporidae). *Mammalian Species* 52: 125-142

576 Bruna T, Hoff KJ, Lomsadze A, Stanke M, Borodovsky M (2021) BRAKER2: automatic eukaryotic
577 genome annotation with GeneMark-EP+ and AUGUSTUS supported by a protein database. *NAR*
578 *Genom Bioinform* 3: lqaa108

579 Carneiro M, Rubin CJ, Di Palma F, Albert FW, Alfoldi J, Martinez Barrio A, Pielberg G, Rafati N,
580 Sayyab S, Turner-Maier J *et al* (2014) Rabbit genome analysis reveals a polygenic basis for
581 phenotypic change during domestication. *Science* 345: 1074-1079

582 Ceballos FC, Joshi PK, Clark DW, Ramsay M, Wilson JF (2018) Runs of homozygosity: windows into
583 population history and trait architecture. *Nat Rev Genet* 19: 220-234

584 Chen SF, Zhou YQ, Chen YR, Gu J (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor.
585 *Bioinformatics* 34: 884-890

586 Cheng H, Concepcion GT, Feng X, Zhang H, Li H (2021) Haplotype-resolved de novo assembly using
587 phased assembly graphs with hifiasm. *Nat Methods* 18: 170-175

588 Crews S, Ojala D, Posakony J, Nishiguchi J, Attardi G (1979) Nucleotide sequence of a region of
589 human mitochondrial DNA containing the precisely identified origin of replication. *Nature* 277:
590 192-198

591 Fazekas J, Grunt TW, Jensen-Jarolim E, Singer J (2017) Long term storage in liquid nitrogen leads
592 to only minor phenotypic and gene expression changes in the mammary carcinoma model cell
593 line BT474. *Oncotarget* 8: 35076-35087

594 Ferreira MS, Jones MR, Callahan CM, Farelo L, Tolesa Z, Suchentrunk F, Boursot P, Mills LS, Alves
595 PC, Good JM *et al* (2021) The Legacy of Recurrent Introgression during the Radiation of Hares.
596 *Syst Biol* 70: 593-607

597 Fickel J, Hauffe HC, Pecchioli E, Soriguer R, Vapa L, Pitra C (2008) Cladogenesis of the European
598 brown hare (*Lepus europaeus* Pallas, 1778). *Eur J Wildlife Res* 54: 495-510

599 Forsyth NR, Elder FF, Shay JW, Wright WE (2005) Lagomorphs (rabbits, pikas and hares) do not
600 use telomere-directed replicative aging in vitro. *Mech Ageing Dev* 126: 685-691

601 Fuste JM, Wanrooij S, Jemt E, Granycome CE, Cluett TJ, Shi Y, Atanassova N, Holt IJ, Gustafsson
602 CM, Falkenberg M (2010) Mitochondrial RNA polymerase is needed for activation of the origin of
603 light-strand DNA replication. *Mol Cell* 37: 67-78

604 Gaertner K, Michell C, Tapanainen R, Goffart S, Saari S, Soininmaki M, Dufour E, Pohjoismaki JLO
605 (2022) Molecular phenotyping uncovers differences in basic housekeeping functions among
606 closely related species of hares (*Lepus* spp., Lagomorpha: Leporidae). *Mol Ecol*

607 Gertz EM, Yu YK, Agarwala R, Schaffer AA, Altschul SF (2006) Composition-based statistics and
608 translated nucleotide searches: improving the TBLASTN module of BLAST. *BMC Biol* 4: 41

609 Ghurye J, Rhie A, Walenz BP, Schmitt A, Selvaraj S, Pop M, Phillippy AM, Koren S (2019) Integrating
610 Hi-C links with assembly graphs for chromosome-scale assembly. *PLoS Comput Biol* 15: e1007273

611 Giska I, Pimenta J, Farelo L, Boursot P, Hacklander K, Jenny H, Reid N, Montgomery WI, Prodohl
612 PA, Alves PC *et al* (2022) The evolutionary pathways for local adaptation in mountain hares. *Mol*
613 *Ecol* 31: 1487-1503

614 Gustavsson I (1972) Mitotic and meiotic chromosomes of the variable hare (*Lepus timidus* L.), the
615 common hare (*Lepus europaeus* Pall.) and their hybrids. *Hereditas* 67: 27-34

616 Holden ME (2005) Family Gliridae. In: *Mammal Species of the World: A Taxonomic and Geographic*
617 *Reference*, Wilson D.E., Reeder D.M. (eds.) pp. 819-841. Johns Hopkins University Press:
618 Baltimore, USA

619 Howe K, Chow W, Collins J, Pelan S, Pointon D-L, Sims Y, Torrance J, Tracey A, Wood J (2021)
620 Significantly improving the quality of genome assemblies through curation. *GigaScience* 10

621 Jaksic FM (2023) Historical ecology and current status of the European hare
622 in South America: a new species in new countries. *Stud Neotrop Fauna E*

623 Januszczak I, Holroyd N (2023) DTOL Tissue and Blood Sampling Standard Operating Procedure:
624 Chordata: Vertebrata (general). *protocolsio* dx.doi.org/10.17504/protocols.io.n92ldmjool5b/v1

625 Juhling F, Putz J, Bernt M, Donath A, Middendorf M, Florentz C, Stadler PF (2012) Improved
626 systematic tRNA gene annotation allows new insights into the evolution of mitochondrial tRNA
627 structures and into the mechanisms of mitochondrial genome rearrangements. *Nucleic Acids Res*
628 40: 2833-2845

629 Juric I, Aeschbacher S, Coop G (2016) The Strength of Selection against Neanderthal Introgression.
630 *PLoS Genet* 12: e1006340

631 Kim D, Paggi JM, Park C, Bennett C, Salzberg SL (2019) Graph-based genome alignment and
632 genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* 37: 907-915
633 Korstanje R, O'Brien PC, Yang F, Rens W, Bosma AA, van Lith HA, van Zutphen LF, Ferguson-Smith
634 MA (1999) Complete homology maps of the rabbit (*Oryctolagus cuniculus*) and human by
635 reciprocal chromosome painting. *Cytogenet Cell Genet* 86: 317-322
636 Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, Jones SJ, Marra MA (2009)
637 Circos: an information aesthetic for comparative genomics. *Genome Res* 19: 1639-1645
638 Lawniczak MKN, Durbin R, Flicek P, Lindblad-Toh K, Wei X, Archibald JM, Baker WJ, Belov K, Blaxter
639 ML, Marques Bonet T *et al* (2022) Standards recommendations for the Earth BioGenome Project.
640 *Proc Natl Acad Sci U S A* 119
641 Levanen R, Kunnasranta M, Pohjoismaki J (2018a) Mitochondrial DNA introgression at the
642 northern edge of the brown hare (*Lepus europaeus*) range. *Ann Zool Fenn* 55: 15-24
643 Levänen R, Pohjoismäki J, Kunnasranta M (2019) Home ranges of semi-urban brown hares (*Lepus*
644 *europaeus*) and mountain hares (*Lepus timidus*) at northern latitudes. *Ann Zool Fenn* 56: 107–
645 120
646 Levanen R, Thulin CG, Spong G, Pohjoismaki JLO (2018b) Widespread introgression of mountain
647 hare genes into Fennoscandian brown hare populations. *Plos One* 13
648 Li D, Liu CM, Luo R, Sadakane K, Lam TW (2015) MEGAHIT: an ultra-fast single-node solution for
649 large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics* 31: 1674-
650 1676
651 Li H (2018) Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34: 3094-
652 3100
653 Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler transform.
654 *Bioinformatics* 25: 1754-1760
655 Lieberman-Aiden E, van Berkum NL, Williams L, Imakaev M, Ragozcy T, Telling A, Amit I, Lajoie BR,
656 Sabo PJ, Dorschner MO *et al* (2009) Comprehensive mapping of long-range interactions reveals
657 folding principles of the human genome. *Science* 326: 289-293
658 Manni M, Berkeley MR, Seppely M, Zdobnov EM (2021) BUSCO: Assessing Genomic Data Quality
659 and Beyond. *Curr Protoc* 1: e323
660 Marques JP, Seixas FA, Farelo L, Callahan CM, Good JM, Montgomery WI, Reid N, Alves PC, Boursot
661 P, Melo-Ferreira J (2020) An Annotated Draft Genome of the Mountain Hare (*Lepus timidus*).
662 *Genome Biol Evol* 12: 3656-3662
663 Mazzoni CJ, Ciofi C, Waterhouse RM (2023) Biodiversity: an atlas of European reference genomes.
664 *Nature* 619: 252
665 Melo-Ferreira J, Vilela J, Fonseca MM, da Fonseca RR, Boursot P, Alves PC (2014) The elusive
666 nature of adaptive mitochondrial DNA evolution of an arctic lineage prone to frequent
667 introgression. *Genome Biol Evol* 6: 886-896
668 Meng G, Li Y, Yang C, Liu S (2019) MitoZ: a toolkit for animal mitochondrial genome assembly,
669 annotation and visualization. *Nucleic Acids Res* 47: e63
670 Mikheenko A, Prjibelski A, Saveliev V, Antipov D, Gurevich A (2018) Versatile genome assembly
671 evaluation with QUAST-LG. *Bioinformatics* 34: i142-i150
672 Minoudi S, Papapetridis I, Karaiskou N, Chatzinikos E, Triantaphyllidis C, Abatzopoulos TJ,
673 Triantafyllidis A (2018) Genetic analyses of brown hare (*Lepus europaeus*) support limited
674 migration and translocation of Greek populations. *Plos One* 13: e0206327

675 Ognev SI (1940) 157. *Lepus (Eulagus) europeus* Pall. (1778) заяц-русак. In: *Mammals of the USSR*
676 *and adjacent countries: rodents (In Russian)*, pp. 112-161. Akademiya Nauk SSSR: Moscow
677 Open2C, Abdennur N, Fudenberg G, Flyamer IM, Galitsyna AA, Goloborodko A, Imakaev M, Venev
678 SV (2023) Pairtools: from sequencing data to chromosome contacts. *bioRxiv*
679 Ou S, Su W, Liao Y, Chougule K, Agda JRA, Hellinga AJ, Lugo CSB, Elliott TA, Ware D, Peterson T *et*
680 *al* (2019) Benchmarking transposable element annotation methods for creation of a streamlined,
681 comprehensive pipeline. *Genome Biol* 20: 275
682 Petrovan S *Lepus europaeus* (European hare) (2013) [DATASET]
683 Pohjoismaki JLO, Michell C, Levanen R, Smith S (2021) Hybridization with mountain hares
684 increases the functional allelic repertoire in brown hares. *Sci Rep* 11: 15771
685 Ranallo-Benavidez TR, Jaron KS, Schatz MC (2020) GenomeScope 2.0 and Smudgeplot for
686 reference-free profiling of polyploid genomes. *Nat Commun* 11: 1432
687 Reid N (2011) European hare (*Lepus europaeus*) invasion ecology: implication for the
688 conservation of the endemic Irish hare (*Lepus timidus hibernicus*). *Biol Invasions* 13: 559-569
689 Reid N, Montgomery WI (2007) Is naturalisation of the brown hare in Ireland a threat to the
690 endemic Irish hare? *Biol Environ* 107b: 129-138
691 Rikala K (1925) Ajojähdilla [flush hunting of hares]. *Metsästys ja Kalastus* 2: 44-48
692 Schai-Braun SC, Schwienbacher S, Smith S, Hackländer K (2023) Coexistence of European hares
693 and Alpine mountain hares in the Alps: what drives the occurrence and frequency of their
694 hybrids? *Journal of Zoology* 320: 214-225
695 Schai-Braun SC, Weber D, Hacklander K (2013) Spring and autumn habitat preferences of active
696 European hares (*Lepus europaeus*) in an agricultural area with low hare density. *Eur J Wildlife Res*
697 59: 387-397
698 Seluanov A, Vaidya A, Gorbunova V (2010) Establishing primary adult fibroblast cultures from
699 rodents. *J Vis Exp*
700 Siivonen L (1972) *Rusakko*. Kustannusosakeyhtiö Otava, Keuruu, Finland
701 Skov L, Coll Macia M, Lucotte EA, Cavassim MIA, Castellano D, Schierup MH, Munch K (2023)
702 Extraordinary selection on the human X chromosome associated with archaic admixture. *Cell*
703 *Genom* 3: 100274
704 Smit AFA, Hubley R, Green P, 2013-2015. RepeatMasker Open-4.0. <http://www.repeatmasker.org>.
705 Smith RK, Jennings NV, Harris S (2005) A quantitative analysis of the abundance and demography
706 of European hares *Lepus europaeus* in relation to habitat type, intensity of agriculture and
707 climate. *Mammal Rev* 35: 1-24
708 Stott P (2003) Use of space by sympatric European hares (*Lepus europaeus*) and European rabbits
709 (*Oryctolagus cuniculus*) in Australia. *Mamm Biol* 68: 317-327
710 Thulin C-G (2003) The distribution of mountain hares (*Lepus timidus*, L. 1758) in Europe: A
711 challenge from brown hares (*L. europaeus*, Pall. 1778)? *Mammal Rev* 33: 29-42
712 Thulin C-G, Tegelström H (2002) Biased geographical distribution of mitochondrial DNA that
713 passed the species barrier from mountain hares to brown hares (genus *Lepus*): an effect of
714 genetic incompatibility and mating behaviour? *J Zool* 258: 299-306
715 Thulin CG, Winiger A, Tallian AG, Kindberg J (2021) Hunting harvest data in Sweden indicate
716 precipitous decline in the native mountain hare subspecies *Lepus timidus sylvaticus* (heath hare).
717 *J Nat Conserv* 64

718 Uliano-Silva M, Ferreira J, Krasheninnikova K, Darwin Tree of Life C, Formenti G, Abueg L, Torrance
719 J, Myers EW, Durbin R, Blaxter M *et al* (2023) MitoHiFi: a python pipeline for mitochondrial
720 genome assembly from PacBio high fidelity reads. *BMC Bioinformatics* 24: 288
721 Vinogradov AE (1998) Genome size and GC-percent in vertebrates as determined by flow
722 cytometry: The triangular relationship. *Cytometry* 31: 100-109
723 Wheeler TJ, Eddy SR (2013) nhmmer: DNA homology search with profile HMMs. *Bioinformatics*
724 29: 2487-2489
725 Wolf JB, Lindell J, Backstrom N (2010) Speciation genetics: current status and evolving
726 approaches. *Philos Trans R Soc Lond B Biol Sci* 365: 1717-1733
727 Zhou C, McCarthy SA, Durbin R (2023) YaHS: yet another Hi-C scaffolding tool. *Bioinformatics* 39
728